

# Efficient Interleukin-21 Production by Optimization of Codon and Signal Peptide in Chinese Hamster Ovarian Cells

Hee Jun Cho<sup>1†</sup>, Byung Moo Oh<sup>1,3†</sup>, Jong-Tae Kim<sup>1</sup>, Jeewon Lim<sup>1,3</sup>, Sang Yoon Park<sup>1</sup>, Yo Sep Hwang<sup>1,3</sup>, Kyoung Eun Baek<sup>1</sup>, Bo-Yeon Kim<sup>2</sup>, Inpyo Choi<sup>1</sup>, and Hee Gu Lee<sup>1,3\*</sup>

<sup>1</sup>Immunotherapy Convergence Research Center, Korea Research Institute of Bioscience and Biotechnology (KRIBB), Daejeon 34141, Republic of Korea

<sup>2</sup>Anticancer Agent Research Center, Korea Research Institute of Bioscience and Biotechnology (KRIBB), Ochang, Cheongju 28116, Republic of Korea

<sup>3</sup>Department of Biomolecular Science, University of Science and Technology (UST), Daejeon 34113, Republic of Korea

Received: November 22, 2018

Revised: November 28, 2018

Accepted: November 29, 2018

First published online  
December 7, 2018

\*Corresponding author

Phone: +82-42-860-4182;

Fax: +82-42-860-4593;

E-mail: hglee@kribb.re.kr

<sup>†</sup>These authors contributed  
equally to this work.

pISSN 1017-7825, eISSN 1738-8872

Copyright© 2019 by  
The Korean Society for Microbiology  
and Biotechnology

Interleukin-21 is a common  $\gamma$ -chain cytokine that controls the immune responses of B cells, T cells, and natural killer cells. Targeting IL-21 to strengthen the immune system is promising for the development of vaccines as well as anti-infection and anti-tumor therapies. However, the practical application of IL-21 is limited by the high production cost. In this study, we improved IL-21 production by codon optimization and selection of appropriate signal peptide in CHO-K1 cells. Codon-optimized or non-optimized human IL-21 was stably transfected into CHO-K1 cells. IL-21 expression was 10-fold higher for codon-optimized than non-optimized IL-21. We fused five different signal peptides to codon-optimized mature IL-21 and evaluated their effect on IL-21 production. The best result (a 3-fold increase) was obtained using a signal peptide derived from human azurocidin. Furthermore, codon-optimized IL-21 containing the azurocidin signal peptide promoted IFN- $\gamma$  secretion and STAT3 phosphorylation in NK-92 cells similar to codon-optimized IL-21 containing original signal peptide. Collectively, these results indicate that codon optimization and azurocidin signal peptides provide an efficient approach for the high-level production of IL-21 as a biopharmaceutical.

**Keywords:** IL-21, codon optimization, signal peptide, NK cells, CHO cells

## Introduction

Interleukin-21 (IL-21) is a pleiotropic cytokine that regulates various immune responses [1]. IL-21 is predominantly produced in CD4<sup>+</sup> T cells and natural killer T (NKT) cells. However, its receptor (IL-21R) is expressed in multiple cellular components of the innate and adaptive immune systems [2]. Therefore, it has potent immune regulatory properties in the context of infection, autoimmune diseases, and cancer [3–5]. IL-21 enhances the proliferation of CD4<sup>+</sup> T cells and regulates the development of IL-17-producing T-helper T cells, which are associated with autoimmune diseases and inflammatory conditions [6, 7]. IL-21 also enhances the proliferation and cytotoxicity of natural killer (NK) cells and promotes their interferon- $\gamma$  secretion [8, 9].

Although recombinant IL-21 (rhIL-21) therapy has been ongoing in preclinical studies and in Phase I and II clinical trials to treat patients with metastatic melanoma and renal cell cancer [10–12], the therapeutic applications of rhIL-21 are limited by the relatively high production cost.

There are several expression systems producing recombinant protein, such as bacteria, fungi, and mammalian cells. Mammalian cells are proper hosts for the production of recombinant proteins, because they are beneficial for post-translational modifications (PTMs). Most marketed recombinant protein pharmaceuticals were produced by Chinese hamster ovary (CHO) cells [13, 14]. Gene optimization and the selection of an appropriate signal peptide can enhance the production of recombinant proteins in mammalian cells. A number of parameters, including codon usage, mRNA

stability and the GC content, and RNA instability motifs and splicing sites, are considered for gene optimization to improve transcription, translation, and folding of recombinant protein [15, 16]. The selection of proper signal peptides is also critical for establishing a manufacturing process for protein production to improve the correct processing and secretion of recombinant proteins through the transport of the translated proteins into the endoplasmic reticulum [17, 18]. Recent studies have demonstrated the effectiveness of different signal peptides on the production of recombinant protein in CHO cells [19–22].

The aim of the current study was to improve the production of IL-21 in the CHO-K1 cell line. IL-21 codon optimization and the selection of a proper signal peptide improved productivity and maintained the biological properties of IL-21.

## Materials and Methods

### Gene Optimization, Synthesis, and Vector Construction

Codon optimization was performed using the Gene Optimization algorithm of Bioneer (Korea). Human IL-21 cDNA (BC066260) and all codon-optimized DNAs were synthesized and cloned into the pGH cloning vector (Bioneer). All genes encoding IL-21 were cloned into pcDNA3.1/zeo(+)-GS (24) at the AflIII/BamHI site.

### Cell Culture and Transfections

CHO-K1 cells were cultured in DMEM (-)Gln (11960-069; Thermo, USA) containing 10% dialyzed fetal bovine serum (dFBS, 26400-044; Thermo) and GS expression medium supplement (GSEM, G9785; Sigma-Aldrich, USA). Cells were transfected with a vector containing mock, non-optimized, and optimized human IL-21 genes using X-tremeGENE 9 DNA Transfection Reagent (06 365 787 001; Roche, Basel, Switzerland) as described previously [23–25]. After transfection,  $2 \times 10^3$  cells were seeded in 96-well plates and selected with DMEM (-)Gln containing 10% dFBS, GSEM, 300  $\mu$ g/ml zeocin (R250-01; Thermo), and 25  $\mu$ M GS System L-Methionine Sulfoximine (Sigma-Aldrich). IL-21 concentrations in the supernatants were measured when the cell pools reached confluence. To construct single cell clones, cell pools high-expressing non-optimized IL-21 or codon-optimized IL-21 were

gradually expanded and subjected to limited dilution. IL-21 production in the respective clonal lines was measured by ELISA. The top three high-producing cell clones were used for further analyses.

### Signal Peptide Optimization

Five different signal peptides were identified by literature search and fused with the codon-optimized mature IL-21 [16–19]. The origin and amino acid sequence of signal peptides are shown in Table 1.

### Quantification of IL-21 and IFN- $\gamma$ by ELISA

IL-21 and IFN- $\gamma$  cytokines were evaluated in the culture supernatant using specific ELISA kits from Invitrogen [26–28]. NK92 cells were seeded in  $5 \times 10^5$ /ml with IL-2-free media for 24 h. Following starvation, NK92 cells were stimulated with the commercial IL-21 expressed in bacteria (rhIL-21), original signal peptide fused-IL-21 (WT/IL-21), or Azurocidin signal peptide-fused IL-21 (Az/IL-21) for 24 h. IFN- $\gamma$  concentration was measured using specific ELISA kits (Invitrogen) according to the manufacturer's instructions.

### Western Blotting


Western blot analysis was performed as described previously, with minor modifications [29, 30]. In brief, cells were lysed in ice-cold lysis buffer (20 mmol/l Tris (pH 8.0), 137 mM NaCl, 10% glycerol, 1% Nonidet P-40, 1 mM phenylmethylsulfonyl fluoride, protease inhibitor mixture (Roche), 1 mM sodium vanadate). Cell lysates were separated by SDS-PAGE and transferred to a polyvinylidene difluoride membrane (Amersham Biosciences, Amersham, UK). Subsequently, membranes were incubated in TBST supplemented with 5% non-fat dry milk and probed with STAT3 (Cell Signalling Technology, USA) and phosphor-STAT3 (Cell Signalling Technology) antibodies at 4°C overnight. The bound antibodies were visualized with a suitable secondary antibody conjugated to horseradish peroxidase using enhanced chemiluminescence reagent (AB Frontier, Korea) [31, 32].

### Statistical Analysis

Data were obtained from at least three independent experiments. All quantitative data are presented as means  $\pm$  standard deviations and were analyzed using Student's *t*-tests.  $P < 0.05$  was considered statistically significant.

**Table 1.** The origin and amino acid sequence of signal peptides.

Name	Signal peptide sequence	Protein	Accession No.	Organism
HSA	MKWVTFISLLFLFSSAYS	Human serum albumin	P02768	<i>Homo sapiens</i>
RSA	MKWVTFLLLLFISGSAFS	Rat serum albumin	P02770	<i>Rattus norvegicus</i>
AZ	MTRLTVLALLAGLLASSRA	Azurocidin preproprotein	NP_001691	<i>Homo sapiens</i>
TPA	MDAMKRGGLCCVLLLCGAVFVSP	Tissue-type plasminogen activator	P00750	<i>Homo sapiens</i>
H7	MEFGLSWVFLVALFRGVQC	Ig heavy chain signal peptide 7		



**Fig. 1.** Evaluation of IL-21 production by gene optimization. (A) CHO-K1 cells were transfected with the vector expressing WT or gene optimized IL-21. Two different CHO-K1-derived cell pools selected and defined were two WT IL21p (cell pools transfected with WT IL-21) and two opti IL-21p (cell pools transfected with opti IL-21).  $1 \times 10^5$  cells were seeded in 6-well plates for 3 days. The concentration of IL-21 in the supernatant was measured by ELISA. (B) Three high-producing cell clones were constructed as described in Materials and Methods.  $1 \times 10^5$  cells were seeded in 6-well plates for 3 days. The concentration of IL-21 in the supernatant was measured by ELISA.

## Results


### Enhanced Production of IL-21 by Gene Optimization

To investigate the effect of gene optimization on IL-21 expression in CHO-K1 cells, synthetic IL-21 genes were assembled based on codon usage in Chinese hamster (*Cricetulus griseus*) as described in Materials and Methods. Amino acid sequences for non-optimized IL-21 (WT IL-21) and optimized IL-21 (opti IL-21) were identical. The sequence of WT IL-21 and opti IL-21 had the identity of 82.4%. The codon optimization index (COI) increased from 0.71 to 0.84. The GC content was also adjusted from 40% to 51%. To investigate the enhancement in productivity by

gene optimization, CHO-K1 cells were transfected with a vector containing WT or opti IL-21. Transfected cells were seeded in 96-well plates and selected. IL-21 concentrations in the culture supernatants were measured when the cell pools reached more than 90% confluence. Two cell pools with high production of WT or opti IL-21 were gradually expanded. The concentration of IL-21 in the supernatant was analyzed by ELISA when cells reached at least 90% confluence. Higher IL-21 concentrations were observed in two different cell pools transfected with optimized IL-21 (38.8 and 35.7 ng/ml) than in two different cell pools transfected with non-optimized IL-21 (13.7 and 13.5 ng/ml) (Fig. 1A). Next, cell pools were subjected to limiting dilution to establish single cell clones. The top three high-producing clones were selected. IL-21 secretion in respective clonal cell lines was analyzed by ELISA. The average yield of opti IL-21 ( $1,495 \pm 192$  ng/ml) was 10-fold higher than that of WT IL-21 ( $159 \pm 24$  ng/ml) (Fig. 1B). These results indicate that gene optimization enhances the production of IL-21 in CHO-K1 cells.

### Enhanced Secretion of IL-21 by the Azurocidin Signal Peptide

Signal peptides play important roles in the secretion of proteins expressed in mammalian cells. To enhance the secretion of IL-21 in CHO-K1 cells, five promising signal peptides were selected by literature search [19–22]. Their origin and amino acid sequences are listed in Table 1. Five different signal peptides were fused to codon-optimized mature IL-21 (Fig. 2A). To evaluate the effects of signal peptides on IL-21 production, CHO-K1 cells were transfected with a vector expressing IL-21 fused to WT or to five different signals. Cell pools were selected as described in the Materials and Methods. IL-21 concentrations in the culture supernatants were analyzed by ELISA when the cells reached confluence. Although IL-21 production was greater using the rat serum albumin signal peptide (RSAP/mIL21) and tissue plasminogen activator signal peptide (tPASP/mIL21) than using the original signal peptide, the greatest productivity was observed for a cell pool expressing azurocidin signal peptide conjugated-IL-21 (AzSP/mIL21) (Fig. 2B). Although the azurocidin signal peptide improved IL-21 productivity in cell pools, the performance in stable cell lines can differ. Therefore, the AzSP/mIL21 cell pool was subjected to limiting dilution to generate stable single cell clones. IL-21 concentrations in the supernatants of twenty clones were measured by ELISA (data not shown). The top two clones with respect to IL-21 production (opti Az/IL21-5 and opti Az/IL21-12) were selected and subjected



**Fig. 2.** Enhanced production of IL-21 by the azurocidin signal peptide.

(A) WT or five different signal peptides were fused to codon-optimized mature IL-21. (B) CHO-K1 cells were transfected with the vector expressing IL-21 fused to WT or five indicated signal peptides. Cell pools were selected as described in Materials and Methods.  $1 \times 10^5$  cells were seeded in 6-well plates for 3 days. The concentration of IL-21 in the supernatant was measured by ELISA. (C) Two high-producing cell clones were constructed as described in Materials and Methods.  $1 \times 10^5$  cells were seeded in 6-well plates for 3 days. The concentration of IL-21 in the supernatant was measured by ELISA.

to ELISA with two clones producing the optimized original signal peptide IL-21 (opti IL21-3 and opti IL21-11). Based on ELISA, opti Az/IL21-5 and opti Az/IL21-12 cell clones resulted in a more than 2.5-fold increase in the IL-21 concentration compared to the concentrations for opti IL21-3 and opti IL21-11 cell clones (Fig. 2C). These data suggest that the azurocidin signal peptide enhances the secretion of IL-21 in CHO-K1 cells.


### Biological Activities of Codon- and Signal Peptide-Optimized IL-21

IL-21 induces the phosphorylation of STAT3 and promotes the secretion of interferon- $\gamma$  (IFN- $\gamma$ ) in NK cells [2, 9]. To verify the biological activity of optimized IL-21, NK-92 cells were stimulated with various concentrations of commercial IL-21 expressed in bacteria (rhIL-21), original signal peptide IL-21 (WT/IL-21), or azurocidin signal peptide IL-21 (Az/IL-21). After 24 h, secreted IFN- $\gamma$  in the culture supernatant was measured by ELISA. WT/IL-21 and Az/IL-21 produced by CHO-K1 cells showed similar induction of the secretion of IFN- $\gamma$  in NK-92 cells (Fig. 3A). However, commercial rhIL-21 produced by bacteria resulted in lower levels of IFN- $\gamma$  production than those for

WT/IL-21 and Az/IL-21 in mammalian cells (Fig. 3A). Furthermore, western blot analysis showed that Az/IL-21 induced strong STAT3 phosphorylation, even at low concentrations, whereas rhIL-21 resulted in a slight increase in the phosphorylation of STAT3 (Fig. 3B). These data suggest that Az/IL-21 has similar biological effects on IFN- $\gamma$  production and STAT3 phosphorylation to those of WT/IL-21 and better activity than those of commercial rhIL-21.

### Discussion

Most bio-therapeutics are produced in established mammalian cell lines, like CHO cells, which ensure correct glycosylation for the modulation of pharmaceutical efficacy in vivo. However, high production costs by low productivity and time requirements by slow proliferation limit the development and use of these systems [33]. Recently, many studies have aimed to increase the production of recombinant proteins in CHO systems by manipulating various factors [34, 35]. Expression vector engineering technologies, including codon optimization of the gene of interest and selection of appropriate signal



**Fig. 3.** IFN- $\gamma$  production and STAT3 phosphorylation by recombinant IL-21 proteins in human NK-92 cells.

(A) NK-92 cells were serum-starved for 18 h and then treated with indicated concentrations of rhIL-21, WT/IL-21, or Az/IL-21 for 24 h. The amount of secreted IFN- $\gamma$  in the culture supernatant was measured by ELISA. Data represent the mean  $\pm$  S.D. of three individual experiments. \*,  $p < 0.05$ , \*\*,  $p < 0.01$ . (B) Serum-starved NK-92 cells were treated with rhIL-21 or Az/IL-21 for 30 min. Cell lysates were immunoblotted with pSTAT3 or STAT3 antibodies.

peptides, are the most convenient methods to enhance productivity [15, 19].

Previous studies have shown that optimization of various factors, including codon usage bias, codon context, GC content, premature poly A, cryptic splice sites, and CpG dinucleotide content [36]. Our results showed that IL-21 productivity was improved 10-fold by gene optimization (Fig. 2B). Many studies have shown that gene optimization is a powerful method to enhance productivity by improving transcription, translation, and protein folding in non-mammalian or mammalian expression systems [37]. The GC content directly regulates the stability of mRNA and indirectly affects the mRNA level [15]. In this study, gene optimization of *IL-21* resulted in an increase in the COI (0.71 to 0.84), which is a measure of codon usage bias, as well as in the GC content (40% to 51%). Therefore, increased production resulting from gene optimization may be attributed the enhancement of both translation and mRNA level.

Signal peptides regulate efficient transport to the ER, appropriate folding, PTM, and the secretion of translated

proteins to the culture supernatant. Several studies suggest that the production of recombinant protein can be improved by the use of alternative signal peptides [19–22]. In this study, we evaluated the efficiency of five signal peptides on the secretion of IL-21. Three signal peptides (derived from rat serum albumin, azurocidin, and tissue plasminogen activator signal peptides) resulted in higher productivity than that of the original signal peptide, whereas two signal peptides (derived from human serum albumin and H7 signal peptide) were similar to the control peptide with respect to productivity. Most notably, the azurocidin signal peptide showed the highest IL-21 production activity (Fig. 2B). Consistent with our results, a previous study used the azurocidin signal peptide to potently express various proteins in mammalian cells [21]. However, another study demonstrated that the human albumin signal peptide shows the highest secretion activity for the production of recombinant protein among various signal peptides [22]. This apparent discrepancy may be attributed to differences in experimental design or systems. Results may vary depending on the cell line or organism. Furthermore, the secretion efficiency can be affected by the amino acid sequence downstream of a signal peptide [38, 39].

The biological activities of recombinant IL-21 produced by CHO-K1 cells were assessed by measuring IFN- $\gamma$  secretion and STAT3 phosphorylation in NK-92 cells (Fig. 3). Codon-optimized Az/IL-21 (IL-21 secreted by the azurocidin-derived signal peptide) had similar activity to that of codon-optimized WT/IL-21 (IL-21 secreted by the original signal peptide). Thus, although the azurocidin-derived signal peptide increases IL-21 production (Fig. 2), it does not affect the biological activity of IL-21 toward NK cells. However, the biological activity of both Az/IL-21 and WT/IL-21 in CHO-K1 increased compared to that of commercial recombinant IL-21 produced in bacteria (Fig. 3). CHO cells regulate PTMs to improve biological activity and reduce the immunological response to the therapeutic recombinant protein. Proper folding of secreted proteins by disulphide bonds, which can be obtained in mammalian cells, is important for their activity [40, 41]. Therefore, these enhanced activities can be explained by PTMs, such as glycosylation, sialylation, and fucosylation, or proper protein folding.

In conclusion, we demonstrated that codon optimization and Azurocidin signal peptide can improve the production efficiency of human IL-21, which has biological activity, in CHO-K1 cells. Our results provide a potential strategy for the production of recombinant IL-21 as a biopharmaceutical.

## Acknowledgments

This research was supported by the National Research Council of Science and Technology (NST) grant by the Korea government (CRC-15-02-KRIBB) and the National Research Foundation of South Korea (NRF) (No. 2018R1A2B6005472).

## Conflict of Interest

HJ Cho, BY Kim, I Choi, and HG Lee have pending patent applications related to this study. Other authors have no conflict of interest to declare.

## References

- Rochman Y, Spolski R, Leonard WJ. 2009. New insights into the regulation of T cells by gamma(c) family cytokines. *Nat. Rev. Immunol.* **9**: 480-490.
- Spolski R and Leonard WJ. 2014. Interleukin-21: a double-edged sword with therapeutic potential. *Nat. Rev. Drug Discov.* **13**: 379-95.
- Asao H, Okuyama C, Kumaki S, Ishii N, Tsuchiya S, Foster D, et al. 2001. Cutting edge: the common  $\gamma$ -chain is an indispensable subunit of the IL-21 receptor complex. *J. Immunol.* **167**: 1-5.
- Kang S, Myoung J. 2017. Host innate immunity against hepatitis E virus and viral evasion mechanisms. *J. Microbiol. Biotechnol.* **27**: 1727-1735.
- Kang S, Myoung J. 2017. Primary lymphocyte infection models for KSHV and its putative tumorigenesis mechanisms in B cell lymphomas. *J. Microbiol.* **55**: 319-329.
- Spolski R, Leonard WJ. 2008. Interleukin-21: basic biology and implications for cancer and autoimmunity. *Annu. Rev. Immunol.* **26**: 57-79.
- Spolski R, Leonard WJ. 2008. The Yin and Yang of interleukin-21 in allergy, autoimmunity and cancer. *Curr. Opin. Immunol.* **20**: 295-301.
- Parrish-Novak J, Dillon SR, Nelson A, Hammond A, Sprecher C, Gross JA, et al. 2000. Interleukin 21 and its receptor are involved in NK cell expansion and regulation of lymphocyte function. *Nature* **408**: 57-63.
- Park YK, Shin DJ, Cho D, Kim SK, Lee JJ, Shin MG et al. 2012. Interleukin-21 increases direct cytotoxicity and IFN- $\gamma$  production of ex vivo expanded NK cells towards breast cancer cells. *Anticancer Res.* **32**: 839-846.
- Davis ID, Skrumsager BK, Cebon J, et al. 2007. An open-label, two-arm, phase I trial of recombinant human interleukin-21 in patients with metastatic melanoma. *Clin. Cancer Res.* **13**: 3630-3636.
- Thompson JA, Curti BD, Redman BG, et al. 2008. Phase I study of recombinant interleukin-21 in patient with metastatic melanoma and renal cell carcinoma. *J. Clin. Oncol.* **26**: 2034-2039.
- Petrella TM, Tozer R, Belanger K, et al. 2012. Interleukin-21 has activity in patients with metastatic melanoma: a phase II study. *J. Clin. Oncol.* **30**: 3396-3401.
- Zhu J. 2012. Mammalian cell protein expression for biopharmaceutical production. *Biotechnol. Adv.* **30**: 1158-1170.
- Walsh G. 2014. Biopharmaceutical benchmarks. *Nat. Biotechnol.* **32**: 992-1000.
- Khan KH. 2013. Gene expression in Mammalian cells and its applications. *Adv. Pharm. Bull.* **3**: 257-263.
- Hung F, Deng L, Ravnkar P, Condon R, Li B, Do L, et al. 2010. mRNA stability and antibody production in CHO cells: improvement through gene optimization. *Biotechnol. J.* **5**: 393-401.
- You M, Yang Y, Zhong C, Chen F, Wang X, Jia T, et al. 2018. Efficient mAb production in CHO cells with optimized signal peptide, codon, and UTR. *Appl. Microbiol. Biotechnol.* doi: 10.1007/s00253-018-8986-5. [Epub ahead of print]
- Jeiranikhameh M, Moshiri F, Keyhan Falasafi S, Zomorodipour A. 2017. Designing signal peptides for efficient periplasmic expression of human growth hormone in *Escherichia coli*. *J. Microbiol. Biotechnol.* **27**: 1999-2009.
- Attallah C, Etcheverrigaray M, Kratje R, Oggero M. 2017. A highly efficient modified human serum albumin signal peptide to secrete proteins in cells derived from different mammalian species. *Protein Expr. Purif.* **132**: 27-33.
- Haryadi R, Ho S, Kok YJ, Pu HX, Zheng L, Pereira NA, et al. 2015. Optimization of heavy chain and light chain signal peptides for high level expression of therapeutic antibodies in CHO cells. *PLoS One* **10**: e0116878.
- Kober L, Zehe C, Bode J. 2013. Optimized signal peptides for the development of high expressing CHO cell lines. *Biotechnol. Bioeng.* **110**: 1164-1173.
- Knappskog S, Ravneberg H, Gjerdrum C, Trösse C, Stern B, Pryme IF. 2007. The level of synthesis and secretion of *Gussia princeps* luciferase in transfected CHO cells is heavily dependent on the choice of signal peptide. *J. Biotechnol.* **128**: 705-715.
- Cho HJ, Kim JT, Lee SJ, Hwang YS, Park SY, Kim BY, et al. 2018. Protein phosphatase 1B dephosphorylates Rho guanine nucleotide dissociation inhibitor 1 and suppresses cancer cell migration and invasion. *Cancer Lett.* **417**: 141-151.
- Shin JS, Ku KB, Jang Y, Yoon YS, Shin D, Kwon OS, et al. 2017. Comparison of anti-influenza virus activity and pharmacokinetics of oseltamivir free base and oseltamivir phosphate. *J. Microbiol.* **55**: 979-983.
- Kim JH, Lee CH, Lee SW. 2016. Hepatitis C virus infection stimulates transforming growth factor-beta1 expression through up-regulating miR-192. *J. Microbiol.* **54**: 520-526.
- Hong S, Yu JW. 2018. Prolonged exposure to lipopolysaccharide induces NLRP3-independent maturation and secretion of

- interleukin (IL)-1 $\beta$  in macrophages. *J. Microbiol. Biotechnol.* **28**: 115-121.
27. Bhardwaj M, Cho HJ, Paul S, Jakhar R, Khan I, Lee SJ *et al.* 2017. Vitexin induces apoptosis by suppressing autophagy in multi-drug resistant colorectal cancer cells. *Oncotarget.* **9**: 3278-3291.
28. Choi DW, Jung SY, Kang J, Nam YD, Lim SI, Kim KT *et al.* 2018. Immune-enhancing effect of nanometric *Lactobacillus plantarum* nF1 (nLp-nF1) in a mouse model of cyclophosphamide-induced immunosuppression. *J. Microbiol. Biotechnol.* **28**: 218-226.
29. Cho HJ, Hwang YS, Yoon J, Lee M, Lee HG, Daar IO. 2018. EphrinB1 promotes cancer cell migration and invasion through the interaction with RhoGDI1. *Oncogene* **37**: 861-872.
30. Yoon J, Hwang YS, Lee M, Sun J, Cho HJ, Knapik L, *et al.* 2018. TBC1d24-ephrinB2 interaction regulates contact inhibition of locomotion in neural crest cell migration. *Nat. Commun.* **9**: 3491.
31. Kim E, Myoung J. 2018. Hepatitis E virus papain-like cysteine protease inhibits type I interferon induction by down-regulating melanoma differentiation-associated gene 5. *J. Microbiol. Biotechnol.* **28**: 1908-1915.
32. Kang S, Choi C, Choi I, Han KN, Rho SW, Choi J, *et al.* 2018. Hepatitis E virus methyltransferase inhibits type I interferon induction by targeting RIG-I. *J. Microbiol. Biotechnol.* **28**: 1554-1562.
33. Lai T, Yang Y, Ng SK. 2013. Advances in Mammalian cell line development technologies for recombinant protein production. *Pharmaceuticals* **6**: 579-603.
34. Schaub J, Clemens C, Schorn P, Hildebrandt T, Rust W, Mennerich D, *et al.* 2010. CHO gene expression profiling in biopharmaceutical process analysis and design. *Biotechnol. Bioeng.* **105**: 431-438.
35. Reinhart D, Damjanovic L, Kaisermayer C, Kunert R. 2015. Benchmarking of commercially available CHO cell culture media for antibody production. *Appl. Microbiol. Biotechnol.* **99**: 4645-4657.
36. Han JH, Choi YS, Kim WJ, Jeon YH, Lee SK, Lee BJ, *et al.* 2010. Codon optimization enhances protein expression of human peptide deformylase in *E. coli*. *Protein Expr. Purif.* **70**: 224-230.
37. Fath S, Bauer AP, Liss M, Spriestersbach A, Maertens B, Hahn P, *et al.* 2011. Multiparameter RNA and codon optimization: a standardized tool to assess and enhance autologous mammalian gene expression. *PLoS One* **6**: e17596.
38. Andrews DW, Perara E, Lesser C, Lingappa VR. 1988. Sequences beyond the cleavage site influence signal peptide function. *J. Biol. Chem.* **263**: 15791-15798.
39. Wiren KM, Potts JT Jr, Kronenberg HM. 1988. Importance of the propeptide sequence of human preproparathyroid hormone for signal sequence function. *J. Biol. Chem.* **263**: 19771-19777.
40. Butler M, Spearman M. 2014. The choice of mammalian cell host and possibilities for glycosylation engineering. *Curr. Opin. Biotechnol.* **30**: 107-112.
41. Zhang P, Chan KF, Haryadi R, Bardor M, Song Z. 2013. CHO glycosylation mutants as potential host cells to produce therapeutic proteins with enhanced efficacy. *Adv. Biochem. Eng. Biotechnol.* **131**: 63-87.