Prevalence and Characterization of Plasmid-Mediated Quinolone Resistance Determinants *qnr* and *aac(6')-Ib-cr* in Ciprofloxacin-Resistant *Escherichia coli* Isolates from Commercial Layer in Korea

Kwang Won Seo^{1,2} and Young Ju Lee^{1*}

¹College of Veterinary Medicine and Zoonoses Research Institute, Kyungpook National University, Daegu 41566, Republic of Korea

²Department of Basic Sciences, College of Veterinary Medicine, Mississippi State University, MS 39762, USA

The prevalence and characterization of plasmid-mediated quinolone resistance (PMQR) determinants in ciprofloxacin-resistant *Escherichia coli* isolated from a Korean commercial layer farm were studied. A total of 45 ciprofloxacin-resistant *E. coli* isolates were recovered and all isolates were multidrug-resistant. Eight isolates have the PMQR genes *aac(6')-Ib-cr*, *qnrS1*, and *qnrB4*, and seven isolates exhibited double amino acid exchange at both *gyrA* and *parC*, and have high fluoroquinolone minimum inhibitory concentrations. Five transconjugants demonstrated transferability of PMQR and β -lactamase genes and similar antimicrobial resistance. Because PMQR genes in isolates from commercial layer chickens could enter the food supply and directly affect humans, control of ciprofloxacin resistance is needed.

Keywords: Antimicrobial resistance, commercial layer chicken, *Escherichia coli*, fluoroquinolones, plasmid-mediated quinolone resistance

Ciprofloxacin (CIP) is an important synthetic antimicrobial agent for treating bacterial infections in both humans and animals [1]. CIP resistance develops when bacteria alter their response(s) to the use of this antimicrobial drug [2]. Recently, a steady increase in the prevalence of CIP-resistant *Escherichia coli* has been reported worldwide [3, 4]. Along with the recent increase in fluoroquinolone resistance, the prevalence of plasmid-mediated quinolone resistance (PMQR) genes has also been increasing in various Enterobacteriaceae worldwide [5]. Although PMQR determinants confer a low level of quinolone resistance on their own, they also facilitate the acquisition of high-level resistance and resistant mutations among susceptible strains [6].

In Korea, the presence of PMQR genes in *E. coli* isolates from broiler poultry or chicken meat has been reported; however, little is known about PMQR genes in *E. coli* from commercial layer farms, which play an important role in the supply of dietary protein through the production of eggs [7]. Unlike broiler chickens, commercial layer hens are raised in the cage system past 70 weeks of age [8]. Their eggs can be contaminated by penetration through the eggshell from the colonized gut, or from contaminated feces during or after oviposition, and by infections originating from the reproductive organs [9, 10]. Especially, contaminated eggs are a common cause of food poisoning in humans and can cause serious illnesses like *Salmonella* infection [11]. In this study, we surveyed genetically characterized CIP-resistant *E. coli* isolates from commercial layer farms in Korea.

Forty-five *E. coli* isolates demonstrating CIP resistance were selected from among 320 *E. coli* samples collected from feces (from chickens 20 weeks of age) and dust in 16 commercial layer farms (62 flocks) during the period between 2015 and 2017. Using the diffusion test, all CIP-resistant *E. coli* isolates were investigated for antimicrobial resistance using drugs from 9 classes applied to 18 antimicrobial discs (BD Biosciences, Sparks, MD, USA). MIC ranging from 0.06 to 512 mg/L to NAL, CIP, and enrofloxacin (ENR) were determined using standard agar dilution methods according to recommendations of the Clinical & Laboratory Standards Institute [12]. Multidrug-resistance (MDR) was defined as acquired non-susceptibility to at least 1 agent in 3 or more antimicrobial categories [13].

Polymerase chain reaction amplification of the PMQR markers *qnrA*, *qnrB*, *qnrC*, *qnrD*, *qnrS*, and *aac(6')-Ib-cr*, and β -lactamase genes *bla*_{CTX-M}, *bla*_{TEM}, *bla*_{SHV}, and *bla*_{OXA}, was performed as previously described [14–16]. The transfer of PMQR genes and β -lactamase genes was performed by conjugation experiments using the broth mating method with sodium azide-resistant *E. coli* J53 as a recipient [17]. Transconjugants were selected on

Received: March 27, 2020
 Accepted: May 14, 2020

First published online:
 May 15, 2020

*Corresponding author
 Phone: +82-53-950-7793
 Fax: +82-505-950-7793
 E-mail: youngju@knu.ac.kr

pISSN 1017-7825
 eISSN 1738-8872

Copyright© 2020 by
 The Korean Society for
 Microbiology and
 Biotechnology

Table 1. Distribution of multidrug-resistance patterns among 45 ciprofloxacin-resistant *E. coli* isolates from commercial layer.

Antimicrobial resistance class pattern ^a	Frequency	Prevalence (%)
Nine of classes	2	4.4
AMGs, BL/BLICs, CEPs, FPIs, FQs, PCNs, PHs, Qs, TETs	2	4.4
Eight of classes	4	8.9
AMGs, CEPs, FPIs, FQs, PCNs, PHs, Qs, TETs	4	8.9
Seven of classes	14	31.1
AMGs, CEPs, FPIs, FQs, PCNs, Qs, TETs	5	11.1
AMGs, FPIs, FQs, PCNs, PHs, Qs, TETs	4	8.9
AMGs, CEPs, FPIs, FQs, PCNs, Qs, TETs	2	4.4
AMGs, CEPs, FQs, PCNs, PHs, Qs, TETs	1	2.2
AMGs, CEPs, PCNs, PHs, Qs, FPIs	1	2.2
CEPs, FPIs, FQs, PCNs, PHs, Qs, TETs	1	2.2
Six of classes	9	20.0
AMGs, CEPs, FQs, PCNs, Qs, TETs	6	13.3
AMGs, FQs, PCNs, PHs, Qs, TETs	2	4.4
AMGs, FPIs, FQs, PCNs, Qs, TETs	1	2.2
Five of classes	7	15.6
AMGs, FQs, PCNs, Qs, TETs	4	8.9
CEPs, FPIs, FQs, PCNs, Qs	1	2.2
CEPs, FPIs, FQs, Qs, TETs	1	2.2
CEPs, FQs, PHs, Qs, TETs	1	2.2
Four of classes	3	6.7
BL/BLICs, CEPs, FQs, Qs	1	2.2
CEPs, FQs, Qs, TETs	1	2.2
FPIs, FQs, PCNs, Qs	1	2.2
Three of classes	6	13.3
CEPs, FQs, Qs	6	13.3
Total	45	100.0

^aAMGs, aminoglycosides; BL/BLICs, β -lactam/ β -lactamase inhibitor combinations; CEPs, cepheids; FPIs, folate pathway inhibitors; FQs, fluoroquinolones; PCNs, penicillins; PHs, phenicols; Qs, quionolones; TETs, tetracyclines.

MacConkey agar (BD Biosciences) plates with sodium azide (100 μ g/ml; Sigma-Aldrich, USA) and ampicillin or tetracycline (100 μ g/ml; Sigma-Aldrich). To identify mutations in the quinolone-resistance determining region, PCR was performed as specified by Rodríguez-Martínez *et al.* (2006) and Vasilaki *et al.* (2008), respectively [18, 19].

Among the 45 CIP-resistant *E. coli* isolates from commercial layer farms, all isolates showed MDR against 3 to 9 classes of antimicrobial agents (Table 1). The resistance rate against 6 and 7 classes was 20.0% and 31.1%, respectively. Although 100% of the isolates showed resistance to quinolones, > 70% of the isolates showed resistance to penicillins, tetracyclines, and aminoglycosides. However, only 6.7% of isolates showed resistance to β -lactam/ β -lactamase inhibitor combinations. The distributions of the 8 PMQR-positive *E. coli* among the 45 CIP resistant-*E. coli* isolates are shown in Table 2. Two *qnr* genes (*qnrS1* and *qnrB4*) were identified in 5 and 1 *E. coli* isolate(s), respectively, while *aac(6')-Ib-cr* genes were detected in 2 *E. coli* isolates. Among the 8 PMQR-positive *E. coli* isolates, 2 β -lactamase genes (*bla*_{CTX-M-15} and *bla*_{TEM-1}) were identified in 2 and 3 *E. coli* isolates, respectively. In the conjugation test, 5 transconjugants demonstrated transferability of PMQR genes and β -lactamase genes and similar antimicrobial resistance (Table 2).

Seven PMQR-positive isolates exhibited double amino acid exchange at both *gyrA* and *parC*, and MICs $\geq$ 16 mg/l of CIP and ENR were observed against these isolates. Furthermore, one PMQR-positive isolate exhibited a single amino acid exchange at *parC*, and MIC $\geq$ 4 mg/l and $\geq$ 8 mg/l of CIP and ENR were observed against this isolate, respectively (Table 2).

In this study, 100% of CIP-resistant *E. coli* isolates showed MDR against 3 to 9 classes of antimicrobial agents. In a previous study, Wasyl *et al.* (2013) reported that 31.0% of MDR *E. coli* was identified from commercial layer farms in the European Union [3]. The difference in MDR between the commercial layer farms in Korea and those of other countries may also be associated with the use of antibiotics in each country. Aalipour *et al.* (2014) reported that antibiotic consumption (per 1 kg of animal products) in the European Union and United States is 21 and 94 mg/kg/year, respectively, while that in Korea has been reported to be 285.7 mg/kg/year [20]. Especially, in Korea, the mass medication of poultry with fluoroquinolones is still permitted, and the sale volume of enrofloxacin is the highest among all antimicrobials used to treat poultry [21]. In addition, the Korea Animal and Plant Quarantine Agency reported that *E. coli* with resistance to CIP in poultry has increased from 37.0% in 2007 to 80.0% in 2015 [22].

Many studies have reported that the prevalence of PMQR genes varies from 2.2% to 57.0% globally, depending on the resistance mechanisms of the animal [23, 24]. In this study, the *qnrB4*, *qnrS1*, and *aac(6')-Ib* genes were identified in 2.2% (1/45), 11.1% (5/45), and 2.2% (1/45) of CIP resistant-*E. coli* isolates, respectively, from a

Table 2. Characteristics of the 8 plasmid-mediated quinolone resistance-positive *E. coli* isolates from commercial layer.

Isolate	Farm	PMQR genes ^a	β-lactamase genes	Antimicrobial resistance pattern ^b	MICs (μg/ml) ^c			QRDR mutations ^d	
					NA	ENR	CIP	<i>gyrA</i>	<i>parC</i>
Donor									
Gi-CC-4	Farm 3	<i>qnrS1</i>	CTX-M-15	AM, CZ, CF, CXM, CTX, CAZ, TE, SXT, G, C	≥512	64	16	S83L, D87N	S80I
Gi-CC-5	Farm 3	<i>qnrS1</i>	CTX-M-15	AM, AMC, CZ, CF, CXM, CTX, CAZ, TE, SXT, G, C	≥512	128	32	S83L, D87N	S80I
Gi-CC-7	Farm 3	<i>qnrS1</i>	-	AM, TE, G	≥512	8	4	WT	S80I
Gi-CC-27	Farm 4	<i>aac(6')-Ib-cr</i>	TEM-1	AM, CTX, TE, SXT, G	≥512	32	16	S83L, D87N	S80I
Gi-CC-28	Farm 4	<i>aac(6')-Ib-cr</i>	TEM-1	AM, CZ, CF, TE, SXT, G	≥512	64	16	S83L, D87N	S80I
Gi-CC-35	Farm 5	<i>qnrB4</i>	-	AM, TE, SXT, G, C	≥512	32	32	S83L, D87N	S80I
Gi-CC-36	Farm 5	<i>qnrS1</i>	-	AM, CZ, CF, TE, SXT, G	≥512	16	16	D87N	S80I
Gi-CC-37	Farm 5	<i>qnrS1</i>	TEM-1	AM, CZ, FOX, TE, G	≥512	32	32	S83L, D87N	S80I
Recipient									
<i>E. coli</i> J53		NT	NT	NT	4	0.06	0.06	NT	NT
Transconjugants									
Gi-CC-4-T		<i>qnrS1</i>	CTX-M-15	AM, CZ, CXM, CTX, CAZ, TE, SXT, G, C	4	0.125	0.25	NT	NT
Gi-CC-5-T		<i>qnrS1</i>	CTX-M-15	AMC, CZ, CF, CTX, CAZ, TE, SXT, G, C	8	0.25	0.125	NT	NT
Gi-CC-28-T		<i>aac(6')-Ib-cr</i>	-	AM, CZ, CF, TE, SXT	16	0.06	0.25	NT	NT
Gi-CC-35-T		<i>qnrB4</i>	-	AM, TE, C	8	0.125	0.06	NT	NT
Gi-CC-37-T		<i>qnrS1</i>	TEM-1	AM, CZ, FOX, G	4	0.125	0.25	NT	NT

^aPMQR, plasmid-mediated quinolone resistance; NT, not tested^bAM, ampicillin; AMC, amoxicillin-clavulanic acid; CZ, cefazolin; CF, cephalothin; FOX, cefoxitin; CXM, cefuroxime; CAZ, ceftazidime; CTX, cefotaxime; C, chloramphenicol; G, gentamicin; SXT, sulfamethoxazole/trimethoprim; TE, tetracycline^cMICs, minimum inhibitory concentrations; NA, nalidixic acid; ENR, enrofloxacin; CIP, ciprofloxacin^dQRDR, quinolone-resistance determining region; WT, wild type

commercial layer farm. The frequency of *qnrS1* and *aac(6')-Ib* was significantly higher than that in other animal studies [*qnrS1* (2.2%) and *aac(6')-Ib* (1.1%)] in Korea, although the frequency of *qnrB4* (0.5%) was low [25]. To establish the association between mutation and the presence of these PMQR determinants, *gyrA* and *parC* were sequenced in PMQR-positive isolates. Among 8 PMQR-positive *E. coli* isolates, 7 exhibited a double amino acid exchange in both *gyrA* and *parC*, and MICs ≥ 16 mg/l of CIP and ENR were observed against these isolates. In this study, the transconjugants expressed similar antimicrobial resistance patterns and revealed the presence of PMQR genes (*qnrS1*, *qnrB4*, and *aac(6')-Ib-cr*) and β -lactamase genes (*bla*_{CTX-M-15} and *bla*_{TEM-1}). This result is also consistent in that these transconjugants have similar antimicrobial resistance patterns and the same antimicrobial resistance genes of the donor strains in the previous study [26]. Therefore, this result suggests wide dissemination of PMQR and β -lactamase genes through plasmids via horizontal transfer, and that poultry can contribute to the transmission of these genes to humans.

To our knowledge, this study was the first to investigate the molecular characteristics of CIP-resistant *E. coli* isolates from a commercial layer farm in Korea. We demonstrated that PMQR genes have a relatively high prevalence (17.8%) in CIP-resistant *E. coli* isolates with related susceptibility to fluoroquinolones, revealing that *qnrS*, *qnrB*, and *aac(6')-Ib* are the most prevalent PMQR genes in *E. coli* isolates from a commercial layer farm. Additionally, our findings suggest that there is a need to emphasize and enforce rational use of antimicrobials, and that regular antimicrobial susceptibility surveillance is essential in commercial layer chicken farming.

Acknowledgments

This work was supported by the Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry and Fisheries (IPET) through the Agriculture, Food and Rural Affairs Research Center Support Program, funded by the Ministry of Agriculture, Food and Rural Affairs (MAFRA) (716002-7).

Conflicts of Interest

The authors have no financial conflicts of interest to declare.

References

- Hopkins KL, Davies RH, Threlfall EJ. 2005. Mechanisms of quinolone resistance in *Escherichia coli* and *Salmonella*: Recent developments. *Int. J. Antimicrob. Agents* 25: 358-373.
- Jacoby GA. 2005. Mechanisms of resistance to quinolones. *Clin. Infect. Dis.* 2: S120-S126.

3. Wasyl D, Hoszowski A, Zając M, Szulowski K. 2013. Antimicrobial resistance in commensal *Escherichia coli* isolated from animals at slaughter. *Front. Microbiol.* **4**: 221.
4. Xu Y, Yu W, Ma Q, Zhou H. 2015. Occurrence of (fluoro)quinolones and (fluoro)quinolone resistance in soil receiving swine manure for 11 years. *Sci. Total Environ.* **530-531**: 191-197.
5. Poirel L, Cattoir V, Nordmann P. 2012. Plasmid-mediated quinolone resistance; interactions between human, animal, and environmental ecologies. *Front. Microbiol.* **3**: 24.
6. Yang H, Chen H, Yang Q, Chen M, Wang H. 2008. High prevalence of plasmid-mediated quinolone resistance genes *qnr* and *aac(6')-Ib-cr* in clinical isolates of Enterobacteriaceae from nine teaching hospitals in China. *Antimicrob. Agents Chemother.* **52**: 4268-4273.
7. Jabir F, Hague MT. 2010. Study on production performance of ISA Brown strain at Krishibid Firm Ltd. Trishal, Mymensingh. *Bangladesh Res. Publ. J.* **3**: 1039-1044.
8. Islam S, Bari MS, Moni SP, Siddique MZF, Uddin MH, Miazi OF. 2016. Phenotypic characteristics of commercial layer strains, ISA brown and hisex brown. *Int. J. Nat. Sci.* **5**: 41-45.
9. Keller LH, Benson CE, Krotec K, Eckroade RJ. 1995. *Salmonella* enteritidis colonization of the reproductive tract and forming and freshly laid eggs of chickens. *Infect. Immun.* **63**: 2443-2449.
10. De Reu K, Grijspeerdt K, Messens W, Heyndrickx M, Uyttendaele M, Debevere J, et al. 2006. Eggshell factors influencing eggshell penetration and whole egg contamination by different bacteria, including *Salmonella* enteritidis. *Int. J. Food Microbiol.* **112**: 253-260.
11. Foley SL, Lynne AM. 2008. Food animal-associated *Salmonella* challenges: pathogenicity and antimicrobial resistance. *J. Anim. Sci.* **86**: E173-E187.
12. CLSI. 2013. Performance standards for antimicrobial susceptibility testing M100-S23. CLSI, Wayne, PA.
13. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. 2012. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin. Microbiol. Infect.* **18**: 268-281.
14. Briñas L, Zarazaga M, Sáenz Y, Ruiz-larrea F, Torres C. 2002. Beta-lactamases in ampicillin-resistant *Escherichia coli* isolates from foods, humans, and healthy animals. *Antimicrob. Agents Chemother.* **46**: 3156-3163.
15. Pitout JDD, Hossain A, Hanson ND. 2004. Phenotypic and molecular detection of CTX-M- β -Lactamases produced by *Escherichia coli* and *Klebsiella* spp. *J. Clin. Microbiol.* **42**: 5715-5721.
16. Yu T, Jiang X, Fu K, Liu B, Xu D, Ji S, et al. 2015. Detection of extended-spectrum β -lactamase and plasmid-mediated quinolone resistance determinants in *Escherichia coli* isolates from retail meat in China. *J. Food Sci.* **80**: M1039-1043.
17. Tamang MD, Nam HM, Jang GC, Kim SR, Chae MH, Jung SC, et al. 2012. Molecular characterization of extended-spectrum- β -lactamase-producing and plasmid-mediated AmpC β -lactamase-producing *Escherichia coli* isolated from stray dogs in South Korea. *Antimicrob. Agents Chemother.* **56**: 2705-2712.
18. Rodríguez-Martínez JM, Velasco C, Pascual A, García I, Martínez-Martínez L. 2006. Correlation of quinolone resistance levels and differences in basal and quinolone-induced expression from three *qnrA*-containing plasmids. *Clin. Microbiol. Infect.* **12**: 440-445.
19. Vasilaki O, Ntokou E, Ikonomidis A, Sofianou D, Frantzidou F, Alexiou-Daniel S, et al. 2008. Emergence of the plasmid-mediated quinolone resistance gene *qnrS1* in *Escherichia coli* isolates in Greece. *Antimicrob. Agents Chemother.* **52**: 2996-2997.
20. Aalipour F, Mirlohi M, Jalali M. 2014. Determination of antibiotic consumption index for animal originated foods produced in animal husbandry in Iran, 2010. *J. Environ. Heal. Sci. Eng.* **12**: 42.
21. Animal and Plant Quarantine Agency (APQA). 2016. National Antimicrobial Resistance Monitoring Program, Gimcheon, Republic of Korea.
22. Animal and Plant Quarantine Agency (QIA). 2015. National Antimicrobial Resistance Monitoring Program, Anyang, Republic of Korea.
23. Kim JH, Cho JK, Kim KS. 2013. Prevalence and characterization of plasmid-mediated quinolone resistance genes in *Salmonella* isolated from poultry in Korea. *Avian. Pathol.* **42**: 221-229.
24. Liu BT, Liao XP, Yang SS, Wang XM, Li LL, Sun J, et al. 2018. Detection of mutations in the *gyrA* and *parC* genes in *Escherichia coli* isolates carrying plasmid-mediated quinolone resistance genes from diseased food-producing animals. *J. Med. Microbiol.* **61**: 1591-1599.
25. Tamang MD, Nam HM, Chae MH, Kim SR, Gurung M, Jang GC, et al. 2012. Prevalence of plasmid-mediated quinolone resistance determinants among *Escherichia coli* isolated from food animals in Korea. *Foodborne Pathog. Dis.* **9**: 1057-1063.
26. Briaes A, Rodríguez-Martínez JM, Velasco C, de Alba PD, Rodríguez-Baño J, Martínez-Martínez L, et al. 2012. Prevalence of plasmid-mediated quinolone resistance determinants *qnr* and *aac(6')-Ib-cr* in *Escherichia coli* and *Klebsiella pneumoniae* producing extended-spectrum β -lactamases in Spain. *Int. J. Antimicrob. Agents.* **39**: 431-434.