

Redoxcitrinin, a Biogenetic Precursor of Citrinin from Marine Isolate of Fungus *Penicillium* sp.

ZHANG, DAHAI^{1,2}, XIANGUO LI¹, JUNG SOOK KANG³, HONG DAE CHOI⁴, JEE H. JUNG⁵, AND BYENG WHA SON^{2*}

¹College of Chemistry & Chemical Engineering, Ocean University of China, Qingdao 266003, P. R. China

²Department of Chemistry, Pukyong National University, Busan 608-737, Korea

³College of Dentistry, Pusan National University, Busan 602-739, Korea

⁴Department of Chemistry, Donggeui University, Busan 614-714, Korea

⁵College of Pharmacy, Pusan National University, Busan 609-735, Korea

Received: December 11, 2006

Accepted: January 24, 2007

Abstract A chemical analysis of the fermentation of the marine-derived fungus *Penicillium* sp. led to the isolation of a biogenetic precursor of citrinin, redoxcitrinin (**1**), together with polyketide mycotoxins, phenol A (**2**), citrinin H2 (**3**), 4-hydroxymellein (**4**), citrinin (**5**), and phenol A acid (**6**). The structures of compounds **1–6** were determined on the basis of physicochemical data analyses. Among them, compounds **1–3** exhibited a potent radical scavenging activity against 1,1-diphenyl-2-picrylhydrazyl (DPPH) with IC₅₀ values of 27.7, 23.4, and 27.2 μM, respectively.

Keywords: Marine-derived fungus, *Penicillium* sp., redoxcitrinin, biogenetic precursor, citrinin, polyketide

Exploitation of the marine environment has been intriguingly successful in recent years in the search for structurally unusual and biologically highly active natural products [4]. To avoid depletion of marine resources and to enable access to large quantities of interesting compounds, there has been a particular interest in the marine microorganisms that are culturable. We have been studying fungi isolated from marine sources for their potential of providing new bioactive compounds [11]. In a combined approach of biological and chemical screening, we are gaining a thorough understanding of the secondary metabolite pattern of the marine-derived fungi. A marine isolate of the fungus *Penicillium* sp. (MFA446) was selected from our screening program for further studies, because of the 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity of its broth extract. We report here the isolation and structure

elucidation of redoxcitrinin (**1**) [3], together with phenol A (**2**) [5, 8, 9], citrinin H2 (**3**) [8, 9], 4-hydroxymellein (**4**) [6], citrinin (**5**) [5], and phenol A acid (**6**) [5] as well as the biological activity of these compounds.

Fungal Isolation and Culture

The fungal strain, *Penicillium* sp., was isolated from the surface of the marine green alga *Ulva pertusa* collected in Bijin Island, Gyeongnam, Korea and identified based on morphological evaluation and fatty acid methyl ester analysis (Korean Culture Center of Microorganism, Seoul, Korea, a similarity index of 0.65). A voucher specimen was deposited at Pukyong National University with the code MFA446. The isolate was cultured for three weeks (static) at 29°C in SWS medium that consisted of soytone (0.1%), soluble starch (1.0%), and seawater (100%) (20× 1 l).

Extraction and Isolation

The mycelium and broth were separated by filtration, and the whole broth was extracted with EtOAc (20 l) to afford crude extract (1.2 g). The broth extract showed mild DPPH scavenging activity, and the active components were purified by assay-guided isolation.

The broth extract (1.2 g) was subjected to Si gel flash chromatography. Elution was performed with *n*-hexane-EtOAc (stepwise, 0–100% EtOAc) to yield five fractions. Fractions 2–5 on medium-pressure liquid chromatography (MPLC) (ODS) by elution with H₂O-MeOH (gradient) afforded crude compounds **1**, **2** and **3**, **4** and **5**, and **6**, respectively, which were further purified by HPLC (YMC, ODS-A) utilizing a 30 min gradient program of 50% to 100% MeOH in H₂O to furnish **1** (5.1 mg), **2** (31.5 mg), **3** (6.7 mg), **4** (8.4 mg), **5** (3.8 mg), and **6** (3.9 mg), respectively.

*Corresponding author

Phone: 82-51-620-6378; Fax: 82-51-628-8147;

E-mail: sonbw@pknu.ac.kr

Redoxcitrinin (**1**): a colorless solid; $[\alpha]_D^{26} +26^\circ$ (*c* 0.2, EtOH); UV (MeOH), λ_{\max} (log ϵ) 301 (6.29), 342 (5.93) (sh) nm; IR (KBr) ν_{\max} 3,392, 1,708, 1,607, 1,418, 1,276, 1,220, 1,160 cm^{-1} ; LREIMS m/z 236 $[\text{M}]^+$ (85), 218 $[\text{M}-\text{H}_2\text{O}]^+$ (70), 203 $[\text{M}-\text{H}_2\text{O}-\text{CH}_3]^+$ (19), 193 $[\text{M}-\text{COCH}_3]^+$ (100), 176 (74), 165 (46), 161 (44), 147 (24), 91 (38); HREIMS m/z 236.1047 $[\text{M}]^+$ (calcd for $\text{C}_{13}\text{H}_{16}\text{O}_4$, 236.1049). See Table 1 for NMR spectral data.

Phenol A (**2**), citrinin H2 (**3**), and phenol A acid (**6**) were obtained as colorless oils, and 4-hydroxymellein (**4**) and citrinin (**5**) were obtained as colorless solids. Compounds **2-6** showed spectral data virtually identical to those reported in the literature [3, 5, 6, 8, 9].

Structural Elucidation and Bioactivity

Citrinin (**5**), a polyketide mycotoxin isolated from fungi, *Penicillium* or *Aspergillus* species, is a well-known contaminant of a number of agricultural products, and it has been demonstrated to possess nephrotoxic activity in addition to a number of other chronic toxic effects [5]. The known compounds **2-6** were identified by a spectroscopic analysis (^1H and ^{13}C NMR, LREIMS, and $[\alpha]_D$) and in comparison to literature data (Fig. 1) [3, 5, 6, 8, 9].

Redoxcitrinin (**1**), $[\alpha]_D^{26} +26^\circ$ (*c* 0.2, EtOH), was obtained as a colorless solid that yielded a molecular formula of $\text{C}_{13}\text{H}_{16}\text{O}_4$ by the HREIMS ($[\text{M}]^+$ m/z 236.1047 (dev -0.2 mmu)) and ^{13}C NMR methods. The IR spectrum of **1** exhibited bands characteristics of hydroxyl ($3,392\text{ cm}^{-1}$), carbonyl ($1,708\text{ cm}^{-1}$), and aryl ($1,607\text{ cm}^{-1}$) functionalities. The ^1H and ^{13}C NMR data for compound **1**, including the results from COSY, DEPT, HMQC, and HMBC experiments, showed the presence of a hexasubstituted benzene ring possessing two hydroxyl, two methyl, an aldehyde, and a

Table 1. NMR spectroscopic data (^1H : 400 MHz; ^{13}C : 100 MHz, $\text{DMSO}-d_6$) for Redoxcitrinin (**1**).

Position	δ_c , mult.	δ_H (<i>J</i> in Hz)	HMBC (H to C)
1	28.1, CH ₃	1.96, s	3
2	208.6, qC		
3	47.1, CH	4.42, q (7.0)	2, 4, 5, 9, 10
4	141.8, qC		
5	116.5, qC		
6	162.0, qC		
7	109.4, qC		
8	162.0, qC		
9	111.4, qC		
10	16.4, CH ₃	1.32, d (7.0)	2, 3, 4
11	11.8, CH ₃	1.98, s	4, 5, 6
12	8.0, CH ₃	2.03, s	6, 7, 8
13	193.9, CH	9.95, s	
8-OH		13.01, s	7, 8, 9

2-butanon-3-yl groups (Table 1). The EIMS spectrum of **1** showed two prominent fragment ions corresponding to the parent molecule with losses of an acetyl and a 2-butanon-3-yl groups [m/z 193 (M^+-COCH_3), 165 ($\text{M}^+-\text{CH}(\text{CH}_3)\text{COCH}_3$)]. Pentasubstituted benzaldehyde was further supported by UV data [301 nm (log ϵ 6.29), 342 (5.93) (sh)].

The connectivity and assignment of the carbon and proton resonances for **1**, which led to the planar structure for this metabolite, were made by interpretation of HMBC data. Diagnostic HMBC correlations, from H₃-1 to C-3, from H-3 to C-2, C-4, C-5, C-9, and C-10, from H₃-10 to C-2, C-3, and C-4, from H₃-11 to C-4, C-5, and C-6, from H₃-12 to C-6, C-7, and C-8, and from 8-OH to C-7, C-8, and C-9, showed the connection of the C3-C4 bond as well as the position of the 5,7-dimethyl, 6,8-dihydroxyl, and 9-aldehyde groups in **1** (Table 1). On the basis of all the foregoing evidence, we propose that the structure of redoxcitrinin is 3-(3,5-dihydroxy-2,4-dimethyl-6-formyl)buten-2-one (**1**). Despite the structural diversity among compounds **1-3**, **5**, and **6**, it is likely that they all derive from common or similar biogenetic intermediates, presumably as part of a polyketide origin [2, 7]. Accordingly, the configuration of C-3 in **1** was supposed to be 3*S* (Fig. 1).

Compound **1** has not been previously described from a natural source. Redoxcitrinin (**1**) has only been reported as a synthetic product obtained during biosynthetic studies of citrinin (**5**) [1, 2, 7]. Compounds **2** and **3** have been reported not only as the secondary metabolites of *Penicillium citrinum* [8], but also as a byproduct of the thermal and alkaline decompositions of citrinin (**5**), respectively [5, 9].

The antioxidant activity was assessed on the basis of the radical scavenging effect on the DPPH free-radical [10-12]. Compounds **1-3** exhibited mild radical scavenging activity against 1,1-diphenyl-2-picrylhydrazyl (DPPH) with IC_{50} values of 27.7, 23.4, and 27.2 μM , respectively, which




Fig. 1. Polyketide metabolites of *Penicillium* sp.

were less active than the positive control, L-ascorbic acid (IC₅₀, 20 μM).

Acknowledgments

This work was supported by a Korea Science and Engineering Foundation (KOSEF) grant funded by the Korea government (MOST) (No. R01-2004-000-10467-0). Mass spectral data were kindly provided by the Korea Basic Science Institute.

REFERENCES

1. Barber, J., R. H. Carter, M. J. Garson, and J. Staunton. 1981. The biosynthesis of citrinin by *Penicillium citrinum*. *J. Chem. Soc. Perkin Trans. 1*, 2577–2583.
2. Barber, J., A. C. Chapman, and T. D. Howard. 1987. The use of sodium [2-²H₃, 1,2-¹³C₂]acetate in determining the biosynthetic origins of hydrogen atoms in fungal metabolites: The biosynthesis of citrinin by *Penicillium citrinum*. *J. Antibiot.* **40**: 245–248.
3. Barber, J. and J. Staunton. 1981. The total synthesis of some deuterium labeled pentaketide derivatives of orsellinic acid. *J. Chem. Soc. Perkin Trans. 1*, 1685–1689.
4. Blunt, J. W., B. R. Copp, M. H. G. Munro, P. T. Northcote, and M. R. Prinsep. 2006. Marine natural products. *Nat. Prod. Rep.* **23**: 26–78.
5. Clark, B. R., R. J. Capon, E. Lacey, S. Tennant, and J. H. Gill. 2006. Citrinin revisited: From monomers to dimers and beyond. *Org. Biomol. Chem.* **4**: 1520–1528, and references cited therein.
6. Cole, R. J., J. H. Moore, N. D. Davis, J. W. Kirksey, and U. L. Diener. 1971. 4-Hydroxymellein: A new metabolite of *Aspergillus ochraceus*. *J. Agr. Food Chem.* **19**: 909–911.
7. Colombo, L., C. Gennari, D. Potenza, C. Scolastico, F. Aragozzini, and C. Merendi. 1981. Biosynthesis of citrinin and synthesis of its biogenetic precursors. *J. Chem. Soc. Perkin Trans. 1*, 2594–2597.
8. Curtis, R. F., C. H. Hassall, and M. Nazar. 1968. The biosynthesis of phenols. Part XV. Some metabolites of *Penicillium citrinum* related to citrinin. *J. Chem. Soc. C* 85–93.
9. Hirota, M., A. B. Menta, K. Yoneyama, and N. Kitabatake. 2002. A major decomposition product, citrinin H₂, from citrinin on heating with moisture. *Biosci. Biotechnol. Biochem.* **66**: 206–210.
10. Kim, H. J., J. H. Kim, C. H. Lee, and H. J. Kwon. 2006. Gentisyl alcohol, an antioxidant from microbial metabolite, induces angiogenesis *in vitro*. *J. Microbiol. Biotechnol.* **16**: 475–479.
11. Li, X., S.-K. Kim, J. S. Kang, H. D. Choi, and B. W. Son. 2006. Radical scavenging hydroxyphenyl ethanoic acid derivatives from a marine-derived fungus. *J. Microbiol. Biotechnol.* **16**: 637–638.
12. Yun, B.-S., I.-K. Lee, J.-P. Kim, and I.-D. Yoo. 2000. Two *p*-terphenyls from mushroom *Paxillus panuoides* with free radical scavenging activity. *J. Microbiol. Biotechnol.* **10**: 233–237.